

University of Groningen

## Temperature stress increases hybrid incompatibilities in the parasitic wasp genus *Nasonia*

Koevoets, T.; van de Zande, L.; Beukeboom, L. W.

*Published in:*  
Journal of Evolutionary Biology

*DOI:*  
[10.1111/j.1420-9101.2011.02424.x](https://doi.org/10.1111/j.1420-9101.2011.02424.x)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2012

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Koevoets, T., van de Zande, L., & Beukeboom, L. W. (2012). Temperature stress increases hybrid incompatibilities in the parasitic wasp genus *Nasonia*. *Journal of Evolutionary Biology*, 25(2), 304-316. <https://doi.org/10.1111/j.1420-9101.2011.02424.x>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

#### Protocol S1:

DNA extraction protocol Koevoets, van de Zande and Beukeboom, Temperature stress increases hybrid incompatibilities in the parasitic wasp genus *Nasonia*.

Adjusted after Hoarau *et al.* (2007) and Whitlock *et al.* (2008).

Males were ground with a sterile grinder in 96-wells titer plates. 100  $\mu$ L digestion buffer (0.5% SDS, 100mM NaCl, 10 mM Tris:HCl pH 8.0, 25 mM EDTA and freshly added 0.2  $\mu$ g/ $\mu$ L proteinase-K) was added to each sample and incubated over-night at 56°C (plate sealed with ThermalSeal). 40  $\mu$ L 6 M NaCl and 140  $\mu$ L CHCl<sub>3</sub> was added and mixed briefly. The titer plates were again sealed with a ThermalSeal and centrifuged in a Qiagen Sigma 4-15C centrifuge for 20 minutes at 3000 rpm. 60  $\mu$ L of the water phase was transferred to a 96-wells fiber plate (Millipore: Multiscreen<sub>HTS</sub>-FB plate), which was pre-wetted with 60  $\mu$ L binding buffer (6 M NaI and 0.1M Na<sub>2</sub>SO<sub>3</sub>, clarified through standard filter paper, then through 0.45  $\mu$ m filter). The plate was incubated at room temperature for 5 minutes, then centrifuged for 15 minutes at 1000 rpm, followed by 10 minutes at 2000 rpm. The flow-through was discarded, the DNA (attached to the filter plate) was washed with 60  $\mu$ L ice-cold wash buffer [4.2 mL of a stock solution (20 mM Tris-HCl (pH 8), 1 mM EDTA, 0.1 M NaCl) diluted with 10.8 mL absolute ethanol per plate] and centrifuged for 10 minutes at 3000 rpm. The flow-through was discarded and the plate dried at room temperature for 30 minutes. Finally, the samples were eluted with 100  $\mu$ L 56°C elution buffer (1 mM Tris-HCl (pH 8) and 0.1 mM EDTA), incubated for 5 minutes at room temperature and the plate was centrifuged for 5 minutes at 1000 rpm and 5 minutes at 2000 rpm, collecting the DNA in clean 96-wells titer plates. In case of failing DNA extraction (2 out of 10 plates using an earlier stage of the protocol, the protocol as described here, should minimize these failures), the remaining aqueous phase after the CHCl<sub>3</sub> step was removed. On this, the regular high salt-chloroform protocol was followed (Maniatis *et al.*, 1982). DNA extracted using the fiber plates was used for PCR amplification undiluted, while DNA extracted using the high salt-chloroform protocol was diluted 10x.

Hoarau G., Coyer J.A., Stam W.T. & Olsen J.L. 2007. A fast and inexpensive DNA extraction/purification protocol for brown macroalgae. *Mol. Ecol. Notes* 7: 191-193.

Maniatis T., Fritsch E.F. & Sambrook J. 1982. *Molecular cloning: a laboratory manual*, 11th edn. Cold spring harbor laboratory press, New York.

Whitlock R., Hipperson H., Mannarelli M. & Burke T. 2008. A high-throughput protocol for extracting high-purity genomic DNA from plants and animals. *Mol. Ecol. Resour.* 8: 736-741.